## IN THE CLAIMS:

- 1. (Original) A method of treating sleep disordered breathing comprising the step of electrical stimulation of nerves to increase muscle tone of upper airway muscles.
- (Original) A method as claimed in claim 1 whereby the afferent nerves are stimulated.
- 3. (Original) The method of claim 2 whereby the site of electrical stimulation is within or adjacent to the genioglossus muscle.
- 4. (Original) The method of claim 2 whereby the site of electrical stimulation is in the vicinity of the hypoglossal motor nucleus or excitatory afferent nerve pathways leading to this structure.
- 5. (Original) The method of claim 1 whereby the electrical stimulation comprises trains of electrical pulses.
- 6. (Original) The method of claim 5 whereby the train length is approximately 10-30 pulses.
- (Original) A method of treating sleep disordered breathing comprising the step of mechanical stimulation of nerves to increase muscle tone of upper airway muscles.
- (Original) The method of claim 7 whereby mechanical stimulation is performed by a piezo electric mechanical element implanted at a site in the vicinity of the upper airway.
- 9. (Original) The method of claim 8 whereby the piezo-electric mechanical element is implanted within or adjacent to the base of the genioglossus muscle.
- 10. (Original) The method of claim 7 whereby the mechanical stimulation is periodic.
- 11. (Original) The method of claim 10 whereby the period is in the order of several seconds of vibration.
- 12. (Original) The method of claim 7 whereby the mechanical vibration occurs at frequencies in the range of 10-50 Hz.

- 13. (Currently Amended) The method of claim 1 [or 7] whereby stimulation is repeated in accordance with the detected state of the airway.
- 14. (Currently Amended) The method of claim 1 [or 7] whereby stimulation is carried out in accordance with a model of Cheyne-Stokes Respiration.
- 15. (Original) Apparatus for treating respiratory disorders comprising a piezoelectric mechanical element, adapted for implant within or adjacent the base of genioglossus muscle and a controller, adapted to elicit vibration of the element via an electrical signal.
- 16. (Original) A method of detecting respiratory disorders comprising the step of measuring a transthoracic impedance changes via implanted electrodes.
- 17. (Original) The method of claim 16 whereby a first electrode is placed in the left sub-pectoral region.
- 18. (Original) The method of claim 16 whereby a second electrode is placed in the right sub-pectoral region.
- 19. (Original) The method of claim 16 whereby the transthoracic impedance is measured by emitting high frequency electrical pulses.
- 20. (Original) The method of claim 19 whereby the frequency of the pulses is high compared to typical respiration or heart rate.
- 21. (Original) The method of claim 19 whereby the frequency of the pulses is approximately 20 Hz.
- 22. (Original) The method of claim 19 whereby the pulses are of approximately 1mA amplitude.
- 23. (Original) The method of claim 19 whereby the pulses are of approximately 15 microsecond duration.
- 24. (Original) The method of claim 19 whereby an impedance signal is compared to a baseline reference.
- 25. (Original) The method of claim 24 whereby the baseline reference is continuously updated.
- 26. (Original) The method of claim 24 whereby the signal having rhythmic variations at a rate of between approximately 6 and 25 per minute is taken as being indicative of normal respiration.

- 27. (Original) The method of claim 24 whereby the signal having a marked reduction in amplitude compared to the reference is taken as being indicative of an obstructive apnea.
- 28. (Original) The method of claim 24 whereby the signal having a first derivative of near zero is taken as being indicative of central apnea.
- (Original) The method of claim 24 whereby the signal having a crescendodecrescendo pattern with a period of approximately 40 to 120 seconds is taken as being indicative of Cheyne-Stokes Respiration.
- 30. (Original) A method of distinguishing open and closed airway apneic events are distinguished by a combination of implanted electrodes and acoustic transducers.
- 31. (Original) A method of treating respiratory disorders as shown and illustrated in Fig. 1
- 32. (NEW) The method of claim 7 whereby stimulation is repeated in accordance with the detected state of the airway.
- 32. (NEW) The method of claim 7 whereby stimulation is carried out in accordance with a model of Cheyne-Stokes Respiration.